

## Demographic changes and genetic losses in populations of a subterranean rodent (*Ctenomys maulinus brunneus*) affected by a natural catastrophe

By. M. H. GALLARDO and NÉLIDA KÖHLER

*Instituto de Ecología y Evolución, Universidad Austral de Chile, Valdivia, Chile*

*Receipt of Ms. 21. 1. 1994*

*Acceptance of Ms. 21. 6. 1994*

### Abstract

Studied the demographic and genetic effects of a volcanic eruption on two local populations (Río Colorado, Las Raíces) of the fossorial rodent *Ctenomys maulinus brunneus*, in the Andes. Our data represent a unique contribution because the pre-existing demographic data and levels of genetic variation were contrasted with the changes monitored afterwards. Comparative census data of the breeding population size before and after the volcanic eruption, revealed a population decline of about 90% in Río Colorado. An electrophoretic survey of 23 presumptive enzyme loci detected a considerable reduction of genetic variability in both populations. In Río Colorado, the proportion of polymorphic loci (P) decreased from 47.8% to 17.4%, and the expected heterozygosity (H) from 8.9% to 2.8%. In Las Raíces, P decreased from 50% to 0% and H from 13.2% to 0%. Although low genetic variability in fossorial mammals is generally assumed to reflect an adaptation to the stable subterranean niche, in some cases it may be merely the result of genetic drift.

### Introduction

Demographic bottlenecks have received significant attention since they have important evolutionary implications for assessing the genetic consequences of reduced population size (NEI and TAJIMA 1981; NEI et al. 1975; CHAKRABORTY and NEI 1977). These estimates also have implications for conservation projects of mammalian species in fragmented habitats (LANDE and BARROWCLOUGH 1987; MARUYAMA and KIMURA 1980; VARVIO et al. 1986) and for speciation events promoted by founder effects (BARTON 1989). Genetic diversity is crucial in an evolutionary sense, and bottlenecks are the quickest means available for losing genetic variation in natural populations through random fluctuations of population size (LEBERG 1992). There is considerable theoretical support for the hypothesis that a natural population passing through a bottleneck should lose genetic variation in direct proportion to the severity of such an event (CHAKRABORTY and NEI 1977; MARUYAMA and FUERST 1985a, b; NEI et al. 1975). Despite the theoretical attention and well developed models predicting bottleneck effects (McCOMMAS and BRYANT 1990), no actual hints of both demographic and genetic data before and after a bottleneck in the wild have yet been reported. Exemplary studies in mammals (BONNELL and SELANDER 1974; HARWOOD and HALL 1990; O'BRIEN and EVERMAN 1988; O'BRIEN et al. 1987; PACKER et al. 1991) have used reduced genetic variation to infer historical bottlenecks due to the absence of previous suitable parameters. To assess the degree of demographic modifications after a volcanic eruption affecting the fossorial rodent *Ctenomys maulinus brunneus* in southern Chile, we present comparative census data. By comparing electrophoretic data from the same 23 presumptive gene loci before and after the bottleneck, we aim to test the hypothesis that bottlenecks lead to predictable decreases in allozyme variation.

## Material and methods

Field studies were conducted in the austral summer 1986–1987, and 1991–1992, at the type locality of *C. maulinus brunneus* (Río Colorado, Malleco province, Chile, 38° 25' S, 71° 32' W; 1,450 m altitude) and in Cordillera Las Raíces, seven km southeast of the topotype locality (Fig. 1). Río Colorado is a flat 4 km<sup>2</sup> high Andean steppe located 3.5 km SW of the Lonquimay volcano. This area sustains a large population of *Ctenomys*, fairly isolated from other local demes by physiographic and vegetational gaps. The Holocene Lonquimay volcano was dormant until December 1988 when its 50-year period of inactivity was interrupted by a 6-months eruptive phase (MORENO and GARDEWEG 1989; BARRIENTOS and ACEVEDO-ARANGUIZ 1992).

Pre-eruption estimates of the number of breeding individuals (N) in Río Colorado were obtained from censuses conducted within four grids differing in size, vegetation type, and plant coverage. All animals were removed from the grids, and their reproductive condition was assessed according to GALLARDO and ANRIQUE (1991).

Site A: a 50 × 70 m grid on consolidated friable soils which had an 80–90% cover of bunchgrass interspersed with annual plants. Site B: two grids (40 × 60 m, and 50 × 50 m) on loose volcanic sediments with a 40–50% cover of annual plants. Site C: a single 60 × 140 m grid on loose volcanic soils which had a 20–25% cover of annual plants. Field observations confirmed a lack of vegetation from sites B and C owing to the deposition of volcanic ash and scoria following the eruption. There was also a lack of *Ctenomys* activity except in site A where a one hectare grid was sampled in the austral summer 1991.

Pre-eruption estimates of allozymic variation obtained from proteins encoded by 23 loci of 51 pre-eruption specimens from Río Colorado and nine from Las Raíces have been previously reported (GALLARDO and KÖHLER 1992; GALLARDO and PALMA 1992). The loci assayed were: isocitrate dehydrogenase (ICD-1, ICD-2, Enzyme Commission No. 1.1.1.42), malate dehydrogenase (MDH-1, MDH-2, 1.1.1.37), glutamate-oxaloacetate transaminase (GOT-1, GOT-2, 2.6.1.1), glycerol-3-




Fig. 1. Map of the study areas depicting the volcanoes (Vn.) that form the Lonquimay Volcanic Front in the Andes of southern Chile

Table 1. Changes in gene frequencies of polymorphic loci in two populations of *Ctenomys m. brunneus* affected by the eruption of volcano Lonquimay, Chile

Pre-eruption data for Río Colorado were obtained from GALLARDO and PALMA (1992), and for Las Raíces, from GALLARDO and KÖHLER (1992)

Locus	Río Colorado		Las Raíces	
	Pre	Post	Pre	Post
Sample size (n)	51	58	9	23
ICD-1				
A	0.020	0.000		
B	0.980	1.000		
MDH-1				
A	0.971	1.000	0.944	1.000
B	0.029	0.000	0.056	0.000
GOT-1				
A	0.010	0.000		
B	0.990	1.000		
GPD				
A	0.961	1.000		
B	0.039	0.000		
Gd				
A			0.000	0.000
B			0.778*	1.000
C			0.222	0.000
GDH				
A			0.000	0.000
B			0.889*	0.000
C			0.111	1.000
LDH-2				
A	0.843*	1.000	0.889*	1.000
B	0.157	0.000	0.111	0.000
XDH-1				
A	0.157	0.034	0.778	1.000
B	0.647*	0.026*	0.000*	0.000
C	0.196	0.940	0.222	0.000
PGM-2				
A	0.814	1.000	0.444	1.000
B	0.186	0.000	0.556	0.000
PGM-3				
A	0.059	0.000	0.000	0.000
B	0.941	1.000	0.944	1.000
C	0.000	0.000	0.056	0.000
PGI				
A	0.951*	1.000	0.000	0.000
B	0.049	0.000	0.944	1.000
C	0.000	0.000	0.056	0.000
PGD				
A	0.078	0.181	0.222	0.000
B	0.902*	0.793*	0.000*	0.000
C	0.020	0.026	0.778	1.000
Mean Polymorphism	47.8 %	17.4 %	50.0 %	0.0 %
Mean N° Allele/locus	1.6	1.3	1.5	1.0
Mean Heterozygosity (direct count)	3.1 %	0.9 %	4.3 %	0.0 %
(Hardy-Weinberg)	8.9 %	2.8 %	13.2 %	0.0 %

\* Genotypes for this locus and population deviate significantly from Hardy-Weinberg expectations.

phosphate dehydrogenase (GPD, 1.1.1.8), glucose 6-phosphate dehydrogenase (Gd, 1.1.1.49), glucose dehydrogenase (GDH, 1.1.1.47), lactate dehydrogenase (LDH-1, LDH-2, 1.1.1.27), xanthine dehydrogenase (XDH), phosphoglucomutase (PGM-2, PGM-3, 2.7.5.1.), glucose isomerase (GPI, 5.3.1.9), phosphogluconate dehydrogenase (PGD, 1.1.1.44), albumin (ALB), transferrin (TRFER), hexokinase (HK, 2.7.1.1), acid phosphatase (ACP-1, ACP-2, 3.1.3.2), and malic enzyme (ME, 1.1.1.40). Fifty eight animals from Río Colorado, and 23 from Las Raíces were live-trapped and screened for the same loci after the eruption. Tissue homogenates, buffer systems, migration conditions and mixtures were prepared according to the methods of SELANDER et al. (1971). Estimates of expected heterozygosity values were compared to the observed numbers and tested for significance by the nonparametric Wilcoxon two-sample test and by a t-test of arcsin transformed genetic data (ARCHIE 1985). All computations of genetic variation were performed with Biosys-1 (SWOFFORD and SELANDER 1989).

## Results

Twenty three animals captured in site A gave an estimate of 66 animals/ha. An additional estimate of 35 animals/ha was generated by pooling density data from the two grids in site B, whereas an estimate of 8.3 animals/ha was obtained from site C. Direct estimates of the breeding population size obtained from census data indicated 1930 animals before the eruption. Based on a density estimate of 20 animals/ha after the eruption in site A, the breeding population size consisted of 168 animals, which corresponds to a 91.3% reduction in population size after the catastrophe. This estimate provided a rough approximation of the effective population size; at best, it was an upper limit, below which  $N_e$  decreases depending on departures from idealized structure (HUSBAND and BARRETT 1992).

There was a severe loss of electrophoretic variation in both populations following this bottleneck episode (Tab. 1). Mean polymorphism was significantly reduced; it dropped from 47.8% to 17.4% in Río Colorado and reached its minimum value in Las Raíces. Eighty percent of previously polymorphic loci in Río Colorado, and all variable loci in Las Raíces became monomorphic after the bottleneck. As theory predicts, the most frequent allele was more likely to become fixed, but the reverse held for loci GDH and PGM-2 in Las Raíces. Similarly, locus XDH-1, although still polymorphic in Río Colorado, changed significantly in frequency after the eruption (Tab. 1).

The difference between the pre- and post-eruption frequency of heterozygotes was highly significant ( $P < 0.0056$ ; Tab. 2). Further indications for rejecting the null hypothesis of equality between initial and final mean heterozygosities were obtained by congruently significant results of angular-transformed data ( $P < 0.004$ ; not shown). No analogous comparisons were conducted in the Las Raíces sample where the final absence of heterozygotes resulted from a generalized monomorphism.

## Discussion

Our estimations of density fall within the range reported for other subterranean mammals (NEVO 1979), implying a direct

Table 2. Observed pre- and post-bottleneck mean heterozygosity values of polymorphic loci in *Ctenomys maulinus brunneus* (Río Colorado)

The non parametric comparison of mean values is based on the Wilcoxon two-sample test

Locus	Pre-bottleneck Heterozygosity	Post-bottleneck Heterozygosity
ICD-1	0.039	0.000
MDH-1	0.058	0.000
GOT-1	0.020	0.000
GPD	0.076	0.000
LDH-2	0.267	0.000
XDH-1	0.523	0.116
PGM-2	0.306	0.000
PGM-3	0.112	0.000
PGI-1	0.094	0.000
PGD	0.182	0.340
HK	0.179	0.144
ACP-2	0.179	0.051
Total	0.170	0.054
Wilcoxon test: Z = -2.7651		P = 0.0056

correlation between number of animals and food supply. Estimates ranging from 3–47 animals/ha in *C. peruanus* (PEARSON 1959) to 218 animals/ha in *C. talarum* (PEARSON et al. 1968) indicate the large variance of demographic attributes in the genus (REIG et al. 1990).

Theoretically, the major genetic consequence of a bottleneck involves a reduction in the number of alleles, because the variants most at risk are those in low frequency (CHAKRABORTY and NEI 1977). This prediction, supported by our data, reached its maximum expression in the Las Raíces sample. Mean heterozygosity was also affected, especially in Las Raíces where all estimates of genetic diversity reached the most extreme minimum values. Although theory predicts a significant reduction of the number of alleles with less than 10 founders, and even fewer pioneers to affect the average heterozygosity (CHAKRABORTY and NEI 1977; MCCOMMAS and BRYANT 1990), significant losses in genetic diversity occurred with a larger number of founders, provided a slow population recovery (LANDE 1987).

In connection with the genetic effects of demographic bottlenecks, deficiencies in sampling may distort the assessment of heterozygosity levels. Although these estimates appear to be more affected by the number of loci screened than by the number of individuals analyzed (GORMAN and RENZI 1979). Considering that the analysis of 8–12 individuals yields, on the average, a heterozygosity estimate within 1% of the value calculated using larger numbers of individuals, sampling biases can be confidently disregarded in our estimations (GORMAN and RENZI 1979). Another possible bias stems from the assumption that losses result from the bottleneck itself (BOILEAU et al. 1992), with no additional genetic cost because a fast demographic recovery is expected. These predictions result in underestimates of the absolute losses, as *Ctenomys* is a k-strategist exhibiting a low intrinsic growth rate (REIG et al. 1990).

Considering the small period of time that has elapsed since the bottleneck, drastic declines in heterozygosity in relation to expected values probably do not result from consanguineous matings, but from a generalized form of inbreeding associated with limited population size (CHESSER and RYMAN 1986). In this respect, panmictic conditions do not hold since considerable levels of intrademic genetic differentiation strongly suggest population subdivision (GALLARDO unpubl. results). Thus, agonistic behaviour and limited dispersal in *Ctenomys* (GALLARDO and ANRIQUE 1991) are not consistent with the assumption of a panmictic unit of 168 individuals, but support an explanation based on the sustained effects of drift acting on disrupted breeding assemblages.

From a conservation viewpoint, bottleneck-flush cycles are major factors contributing to species vulnerability (O'BRIEN and EVERMAN 1988). Limited population sizes threaten demic persistence when a threshold density value defining mating success is reached (LANDE 1987, 1988). Besides, the genetic uniformity attained through repeated bottleneck events also affects demic survival by inbreeding depression (PACKER et al. 1991; WILDT et al. 1987), or by a homogeneous immunological response to environmental disturbances (O'BRIEN et al. 1985).

Although current interpretations of genic and chromosomal variation patterns in *Ctenomys* emphasize a causal relationship with spatial components (REIG et al. 1990), sustained vulcanism and the tectonically-derived seismic activity in the Andes heighten a component of temporal stochasticity. In this line, previous eruptions of the Lonquimay volcano took place in 1940, 1887 and 1853 (MORENO and GARDEWEG 1989). Assuming similar environmental effects, population survivorship, and a one-year generation time (GALLARDO and ANRIQUE 1991) the genetic pool of these, and probably other local populations has been altered in generations 135, 101, and 48 before the present.

While contrasting with the multiple karyotypic forms observed in the low land *Ctenomys* species (GALLARDO 1991), the karyotypic uniformity reported for species from the Andes (GALLARDO 1979) conflicts with the optimal conditions for chromosomal differentiation that stem from limited population size (LANDE 1979, 1985; WRIGHT 1941).

Apparently, chromosomal conservatism may be better explained by recurrent vulcanism for when extinctions and recolonizations are frequent, subpopulation divergence is prevented (MARUYAMA and KIMURA 1980) and new colonies are likely to descend from a single ancestral deme (WRIGHT 1941). Considering that the activity of the nearby volcanoes Antuco, Llama and Villarrica total 36 eruptive events since 1640 (VEBLEN 1985), the long-term karyotypic uniformity in Andean *Ctenomys* populations appears to be coupled to the structure of the environment. Furthermore, low levels of genetic variability attained by recurrent catastrophes may imitate an adaptive response to the stable subterranean niche (NEVO 1979, 1990) although no selective pressure can counterbalance the effects of drift when populations recover from precarious levels of genetic variability (MARUYAMA and FUERST 1985a, b).

### Acknowledgements

This research was supported by grant FN 92/0178 and DID UACH grant S-94-29 to MHG.

### Zusammenfassung

#### *Demographische Veränderungen und Verluste von genetischer Variabilität bei Populationen des grabenden Nagers (Ctenomys maulinus brunneus) als Folge einer Naturkatastrophe*

Es wurden die demographischen und genetischen Auswirkungen eines Vulkanausbruchs auf zwei lokale Populationen (Río Colorado, Las Raíces) des grabenden Nagers *Ctenomys maulinus brunneus* in den Anden untersucht. Ein Vergleich der Populationsgrößen vor und nach dem Vulkanausbruch auf der Grundlage von Zählungsergebnissen in Río Colorado ergab einen Rückgang der Individuenzahlen um etwa 90%. Die elektrophoretische Untersuchung von 23 Enzymloci zeigte in beiden Beständen eine starke Reduktion der genetischen Variabilität. In Río Colorado sanken die Polymorphierate (P) von 47,8% auf 17,4% und der durchschnittliche erwartete Heterozygotiegrad von 8,9% auf 2,8%. In Las Raíces sanken P von 50% auf 0% und H von 13,2% auf 0%. Unsere Daten liefern einen Beitrag zu den wenigen Fällen, in welchen bei natürlichen Populationen Verluste von genetischer Variabilität direkt einem Engpaß in der Populationsgröße zugeordnet werden können. Obwohl die geringe genetische Variation bei grabenden Nagern in der Regel als Anpassung an die stabilen Lebensbedingungen der unterirdischen Nische zugeschrieben wird, kann sie in einigen Fällen lediglich auf genetische Drift zurückzuführen sein.

### Literature

- ARCHIE, J. W. (1985): Statistical analysis of heterozygosity data: independent sample comparisons. *Evolution* **39**, 623–637.
- BARRIENTOS, S. E.; ACEVEDO-ARANGUIZ, P. S. (1992): Seismological aspects of the 1988–1989 Lonquimay (Chile) volcanic eruption. *J. Volcan. Geothermal Res.* **53**, 73–87.
- BARTON, N. H. (1989): Founder effect speciation. In: *Speciation and its consequences*. Ed. by D. OTTE and J. A. ENDLER. Massachusetts: Sinauer Ass. Pp. 229–256.
- BOILEAU, M. G.; HEBERT, P. D. N.; SCHWARTZ, S. S. (1992): Non-equilibrium gene frequency divergence: persistence founder effects in natural populations. *J. Evol. Biol.* **5**, 25–39.
- BONNELL, M. L.; SELANDER, R. K. (1974): Elephant seals: genetic variation and near extinction. *Science* **184**, 908–909.
- CHAKRABORTY, R.; NEI, M. (1977): Bottleneck effects on average heterozygosity and genetic distance with the stepwise mutation model. *Evolution* **31**, 347–356.
- CHESSER, R. K.; RYMAN, N. (1986): Inbreeding strategy in subdivided populations. *Evolution* **40**, 616–624.
- GALLARDO, M. H. (1979): Las especies chilenas de *Ctenomys*. I. Estabilidad cariotípica. *Arch. Biol. Med. Exper.* **12**, 71–82.
- (1991): Karyotypic evolution in *Ctenomys* (Rodentia, Ctenomyidae). *J. Mammalogy* **72**, 11–21.
- GALLARDO, M. H.; ANRIQUE, J. (1991): Populational parameters and burrow systems in *Ctenomys maulinus brunneus* (Rodentia, Ctenomyidae). *Med. Ambiente* **11**, 48–53.
- GALLARDO, M. H.; KÖHLER, N. (1992): Genetic divergence in *Ctenomys* (Rodentia, Ctenomyidae) from the Andes of Chile. *J. Mammalogy* **72**, 99–105.
- GALLARDO, M. H.; PALMA, R. E. (1992): Intra- and interspecific genetic variability in *Ctenomys* (Rodentia, Ctenomyidae). *Biochem. Syst. Ecol.* **20**, 523–534.
- GORMAN, G. C.; RENZI, J. (1979): Genetic distance and heterozygosity estimates in electrophoretic studies: effects of sample size. *Copeia* **2**, 242–249.

- HARWOOD, J.; HALL, A. (1990): Mass mortality in marine mammals: its implications for population dynamics and genetics. *Trends Ecol. Evol.* 5, 254–257.
- HUSBAND, B. C.; BARRETT, S. C. H. (1992): Effective population size and genetic drift in *Eichhornia paniculata*. *Evolution* 46, 1875–1890.
- LANDE, R. (1979): Effective deme sizes during long-term evolution estimated from rates of chromosomal rearrangements. *Evolution* 33, 234–251.
- (1985): The fixation of chromosomal rearrangements in a subdivided population with local extinction and colonization. *Heredity* 54, 323–332.
- (1987): Extinction thresholds in demographic models of territorial populations. *Amer. Naturalist* 130, 624–635.
- (1988): Genetics and demography in biological conservation. *Science* 241, 1455–1460.
- LANDE, R.; BARROWCLOUGH, G. F. (1987): Effective population size, genetic variation, and their use in population management. In: *Viable populations for conservation*. Ed. by M. E. SOULÉ. New York: Cambridge Univ. Press. Pp. 87–123.
- LEBERG, P. (1992): Effects of population bottlenecks on genetic diversity as measured by allozyme electrophoresis. *Evolution* 46, 477–494.
- MARUYAMA, T.; FUERST, P. A. (1985a): Population bottlenecks and nonequilibrium models in population genetics. III. Genic homozygosity in populations which experience periodic bottlenecks. *Genetics* 111, 691–703.
- (1985b): Population bottlenecks and nonequilibrium models in population genetics. II. Number of alleles in a small population that was formed by a recent bottleneck. *Genetics* 111, 675–689.
- MARUYAMA, T.; KIMURA, M. (1980): Genetic variability and effective population size when local extinction and recolonization of subpopulations are frequent. *Proc. Natl. Acad. Sci. USA* 77, 6710–6714.
- MCCOMMAS, S. A.; BRYANT, E. H. (1990): Loss of electrophoretic variation in serially bottlenecked populations. *Heredity* 64, 315–321.
- MORENO, H.; GARDEWEG, M. C. (1989): La erupción reciente en el complejo volcánico Lonquimay (Diciembre 1988). *Andes del Sur. Rev. Geol. Chile* 16, 93–117.
- NEI, M.; MARUYAMA, T.; CHAKRABORTY, R. (1975): The bottleneck effect and genetic variability in populations. *Evolution* 29, 1–10.
- NEI, M.; TAJIMA, F. (1981): Genetic drift and estimation of effective population size. *Genetics* 98, 625–640.
- NEVO, E. (1979): Adaptive convergence and divergence of subterranean mammals. *Ann. Rev. Ecol. Syst.* 10, 269–368.
- (1990): Genetic diversity and its ecological correlates in nature: comparisons between subterranean, fossorial, and aboveground small mammals. In: *Evolution of subterranean mammals at the organismal and molecular levels*. Ed. by E. NEVO and O. A. REIG. New York: A. R. Liss. Pp. 347–366.
- O'BRIEN, S. J.; EVERMAN, J. F. (1988): Interactive influence of infectious disease and genetic diversity in natural populations. *Trends Ecol. Evol.* 3, 254–259.
- O'BRIEN, S. J.; ROELKE, M. E.; MARKER, L.; NEWMAN, A.; WINKLER, C. A.; MELTZER, D.; COLLY, L.; EVERMANN, J. F.; BUSH, M.; WILDT, D. E. (1985): Genetic basis for species vulnerability in the cheetah. *Science* 227, 1428–1435.
- O'BRIEN, S. J.; WILDT, D. E.; BUSH, M.; CARO, T. M.; FITZGIBBON, C.; AGGUNDEY, I.; LEAKEY, R. E. (1987): East African cheetahs: evidence for two population bottlenecks. *Proc. Natl. Acad. Sci. USA* 84, 508–511.
- PACKER, C.; PUSEY, A. E.; ROWLEY, H.; GILBERT, D. A.; MARTENSON, J.; O'BRIEN, S. J. (1991): Case study of a population bottleneck: lions of the Ngorongoro crater. *Conserv. Biol.* 5, 229–237.
- PEARSON, O. P. (1959): Biology of subterranean rodents, *Ctenomys*, in Peru. *Mem. Mus. Hist. Nat. "Javier Prado"* 9, 1–56.
- PEARSON, O. P.; BINSTEIN, N.; BOIRY, L.; BUSH, M. C.; DI PACE, M.; GALLOPIN, G.; PENCHASZADEH, P.; PIANATANIDA, M. (1968): Estructura social, distribución espacial y composición por edades de una población de tuco-tucos (*Ctenomys talarum*). *Invest. Zool. Chile* 13, 47–80.
- REIG, O. A.; BUSH, C.; ORTELLS, M. O.; CONTERAS, J. R. (1990): An overview of evolution, systematics, population biology, cytogenetics, molecular biology, and speciation in *Ctenomys*. In: *Evolution of subterranean mammals at the organismal and molecular levels*. Ed. by E. NEVO and O. A. REIG. New York: A. R. Liss. Pp. 71–96.
- SELANDER, R. K.; SMITH, M. H.; YANG, S. Y.; JOHNSON, W. E.; GENTRY, J. B. (1971): Biochemical polymorphism and systematics in the genus *Peromyscus*. I. Variation in the old-field mouse *Peromyscus polionotus*. *Univ. Texas Publ.* 7103, 49–90.
- SWOFFORD, D. L.; SELANDER, R. (1989): Biosys-1. A computer program for the analysis of allelic variation in population genetics and biochemical systematics. Illinois Natural History Survey.
- VARVIO, S. L.; CHAKRABORTY, R.; NEI, M. (1986): Genetic variation in subdivided populations and conservation genetics. *Heredity* 67, 189–198.
- VEBLEN, T. T. (1985): Stand dynamics in Chilean *Nothofagus* forests. In: *The ecology of natural*

- disturbance and patch dynamics. Ed. by S. T. A. PICKETT and P. S. WHITE. London: Acad. Press. Pp. 35-51.
- WILDT, D. E.; BUSH, M.; GOODGROWE, K. L.; PACKER, C.; POSEY, A. E.; BROWN, J. L.; JOSLIN, P.; O'BRIEN, S. J. (1987): Reproductive and genetic consequences of founding isolated lion populations. *Science* **329**, 328-330.
- WRIGHT, S. (1931): Evolution in mendelian populations. *Genetics* **16**, 97-159.
- (1941): On the probability of fixation of a reciprocal translocation. *Amer. Naturalist* **75**, 513-522.
- Authors' address:* MILTON H. GALLARDO and NÉLIDA KÖHLER, Instituto de Ecología y Evolución, Universidad Austral de Chile, Casilla 567, Valdivia, Chile